

Theoretical and Phenomenological Aspects of $G\chi PT^{*\dagger}$

Jan Stern^{a‡}

^aDivision de Physique Théorique, Institut de Physique Nucléaire,
91406 Orsay, France

Consequences of the alternative mechanism of spontaneous breaking of chiral symmetry ($SB\chi S$) without a formation of a large quark antiquark condensate is reviewed. Emphasis is put on the resulting picture of light quark masses.

1. THEORETICAL ASPECTS OF $SB\chi S$ IN QCD

The spontaneous breaking of chiral symmetry $SU_L(N_f) \times SU_R(N_f) \times U_V(1)$, (associated with N_f massless flavours), down to the diagonal subgroup $U_V(N_f)$ is one of our theoretical certitudes in QCD [1,2]. It is equivalent to the statement that in the chiral limit $m \rightarrow 0$, the left-right correlator

$$F^2 \delta^{ij} = \lim_{m \rightarrow 0} i \eta^{\mu\nu} \int dx < \Omega | T J_{\mu L}^i(x) J_{\nu R}^j(0) | \Omega > \neq 0, \quad (1)$$

where $J_{\mu L}$, ($J_{\mu R}$) are Noether currents generating left (right) chiral rotations, remains non-zero. (F is nothing but the pion decay constant $F_\pi = 92.4$ MeV at $m = 0$). Indeed, Eq. (1) encompasses both essential features of $SB\chi S$: **i**) The asymmetry of the vacuum and, **ii**) the existence of $N_f^2 - 1$ massless Goldstone bosons coupled to the Noether currents with the strength F . In other words, F is an **order parameter**, whose non-vanishing is not only a sufficient but also a necessary condition of $SB\chi S$. There are, of course, many other order parameters, such as the $q\bar{q}$ condensate ($N_f = 2$, $m_u = m_d = m$)

$$\lim_{m \rightarrow 0} < \Omega | \bar{u}u | \Omega > = \lim_{m \rightarrow 0} < \Omega | \bar{d}d | \Omega > = -F^2 B, \quad (2)$$

*IPNO/TH 97-29

[†]Invited talk at the workshop QCD97, Montpellier, France, July 1997

[‡]Laboratoire des Universités Paris XI et Paris VI associé au CNRS.

playing a different role than the pion decay constant F : $B \neq 0$ is **not** a necessary consequence of $SB\chi S$, rather, the magnitude of B reflects the nature of the chiral order in the QCD vacuum. Quark condensate is analogous to the spontaneous magnetization $\vec{m}$ of spin systems, reflecting the type of the magnetic order in the ground state : $\vec{m} \neq 0$ for a ferromagnet, but $\vec{m} = 0$ for an antiferromagnet.

The non-equivalence of quark condensation and of $SB\chi S$ may be illustrated treating the Euclidean QCD as a disordered system[2]. Indeed, the mechanism of $SB\chi S$ can be analyzed in terms of basic characteristics of lowest eigenstates of (Euclidean) Dirac operator in a random gluonic background : density and localization. A $q\bar{q}$ condensate is formed, provided the density of such states is large enough : the average number of eigenvalues in a given small neighbourhood of zero should be proportional to the volume V [3]. On the other hand, F_π - Eq. (1) - remains non zero, i.e. $SB\chi S$ takes place and Goldstone bosons are formed under somewhat weaker conditions. $F \neq 0$ can actually be realized **either** with a **large** density of **localized** states, **or** with a **lower** density (number of small eigenvalues $\sim \sqrt{V}$) of **delocalized** states. In the first case quarks do condense ($B \neq 0$), whereas in the second case $B \rightarrow 0$ in the thermodynamic limit $V \rightarrow \infty$ [4].

The question of the size of the quark condensate is of theoretical nature : there is no known way how to decide a priori which of the two mechanisms summarized above takes actually place

and, even if $B \neq 0$, how close are we to the pure $B = 0$ scenario. In practice, one has to decide whether B is as large as ~ 2 GeV, as suggested by lattice simulations[5], or as small as $B \sim 100$ MeV as it results from a recent attempt to calculate directly in the continuum within the framework of variationally improved perturbation theory[6]. The decision will likely come from new low-energy high precision experiments[8,9].

2. GENERALIZED CHIRAL PERTURBATION THEORY

In Nature, the $\overline{MS}$ running quark masses m_u, m_d, m_s (renormalized at the QCD scale $\nu = 1$ GeV) appear small compared to the typical mass-scale of the bound states $(\rho, N, \dots)$, $\Lambda_H \simeq 1$ GeV. Low energy observables can be expanded in powers of m/Λ_H and in powers of external momenta p/Λ_H . The standard version of χPT assumes, in addition, that the deviation Δ from the GOR relation[7]

$$M_\pi^2 = 2mB + \Delta \quad (3)$$

can be treated as a small perturbation : $S\chi PT$ [10] is a simultaneous expansion in powers of m_q/Λ_H **and** of $\epsilon = \Delta/M_\pi^2$. If B vanishes, $\epsilon = 1$ and the latter expansion breaks down. For nonzero but sufficiently small B , this expansion could still break down, unless quark masses are mathematically small. $G\chi PT$ [11,12] still expands in powers of m/Λ_H avoiding however any expansion in powers of $\epsilon = \Delta/M_\pi^2$. Technically, this is achieved by a different chiral power counting[11] :

$$\begin{aligned} S\chi PT : \quad m_q &= O(p^2) , \quad B = O(1) \\ G\chi PT : \quad m_q &= O(p) , \quad B = O(p) . \end{aligned} \quad (4)$$

In $G\chi PT$, the first term in the expansion (3) of M_π^2 is suppressed, and this is why $2mB$ and Δ could be of a comparable size.

3. EXPANSION OF M_π^2

It is instructive to consider in more details the expansion ($2m = m_u + m_d$)

$$\frac{1}{F^2} F_\pi^2 M_\pi^2 = 2mB +$$

$$+ 4m^2 \left\{ A(\mu) - \frac{3B^2}{32\pi^2 F^2} \ln \frac{M_\pi^2}{\mu^2} \right\} + O(m^3) \quad (5)$$

which, as it stands, is independent of the chiral counting (4). The point we would like to make is that the constant $A(\mu)$ (for $\mu \sim \Lambda_H$) is insensitive to the Goldstone-boson sector of the theory, in particular, to the size of B . This can be seen from the definition of the constant $A(\mu)$ in terms of the two point function of scalar-isoscalar and pseudoscalar-isovector quark densities $S^0(x)$ and $P^i(x)$. For $m_u = m_d = 0$, one has [10]

$$\begin{aligned} & \frac{i}{F^2} \int dx e^{ipx} \langle 0 | T \{ \delta^{ij} S^0(x) S^0(0) - \\ & \quad - P^i(x) P^j(0) \} | 0 \rangle = \\ & = \delta^{ij} \left\{ \frac{B^2}{p^2} + \frac{3}{32\pi^2} \frac{B^2}{F^2} \left(\ln \frac{\mu^2}{-p^2} + 1 \right) + A(\mu) \right. \\ & \quad \left. + O(p^2) \right\} . \end{aligned} \quad (6)$$

The first term on the r.h.s. is the GB-pole, the second term results from the GB-loop, whereas $A(\mu)$ collects contributions from exchanges of non-Goldstone particles : $\sigma, \pi' \dots$. It is hard to believe that the latter would be significantly affected by the mechanism of $SB\chi S$. Comparing Eqs. (5) and (6) with the corresponding $S\chi PT$ expressions[10], one finds

$$A(\mu) = \frac{2B^2}{F^2} [l_3^r(\mu) + l_4^r(\mu)] . \quad (7)$$

This allows one to estimate $A(\mu)$ in the standard case of a large condensate : Taking e.g. $B \simeq 1.6$ GeV (corresponding to $m \simeq 6.1$ MeV) and using the standard values [10], $l_3^r(M_\rho) = 0.8 \times 10^{-3}$, $l_4^r(M_\rho) = 5.6 \times 10^{-3}$, one finds

$$A(M_\rho) \simeq 4 . \quad (8)$$

This value should not be considerably modified in $G\chi PT$. Consequently, if B drops out, the low energy constant $l_3 + l_4$ of $S\chi PT$ should blow up, signaling the inadequacy of the standard expansion, rather than a pathology. If B is small enough to justify the $G\chi PT$ counting (4), the expansion (5) can be rewritten as

$$M_\pi^2 = 2mB + 4m^2 A + O(p^3) . \quad (9)$$

The first two terms are now of the same order $O(p^2)$. The NLO $O(p^3)$ correction includes the

factor $F_\pi^2/F^2 - 1$ as well as a part of $O(m^3)$ contributions in Eq. (5). Finally, the chiral logarithms proportional to $m^2 B^2$ are now suppressed : they are relegated to the order $O(p^4)$. The linear and quadratic terms in Eq. (9) are of comparable size for

$$m \sim m_0 = \frac{B}{2A} . \quad (10)$$

For $B \sim 100$ MeV, $A \sim 4$, m_0 can, indeed, be so small to make problematic any expansion in powers of m/m_0 , in particular, the $S\chi PT$.

4. LOW-ENERGY $\pi\pi$ SCATTERING

The low-energy $\pi\pi$ scattering amplitude is particularly sensitive to the size of the $\bar{q}q$ condensate[11,13]. It's $S\chi PT$ expansion[14] makes appear the low energy constant l_3 and not just the combination $B^2 l_3$ as in Eqs. (5) and (6). It is instructive to look at the low energy parameter α (defined including the $O(p^6)$ accuracy in [13]), which is closely related to the scattering amplitude $A(s, t, u)$ at the symmetric point $s = t = u = \frac{4}{3}M_\pi^2$:

$$A\left(\frac{4}{3}M_\pi^2, \frac{4}{3}M_\pi^2, \frac{4}{3}M_\pi^2\right) = \frac{\alpha}{3F_\pi^2} M_\pi^2 + \dots \quad (11)$$

The parameter α turns out to be more sensitive to the magnitude of B than, say, the S-wave scattering lengths : At the leading order of $G\chi PT$, $\alpha = 1$ in the case of a large condensate, whereas $\alpha \simeq 4$ for $B = 0$. In addition, the parameter α receives rather moderate corrections from higher orders of χPT . Nowadays, the $\pi\pi$ scattering amplitude is analyzed up to and including two-loop accuracy both in the standard[15] and in the generalized[13] frameworks. Within the $S\chi PT$ one gets for instance

$$\alpha = 1 + \left(6l_3^r + 2l_4^r - \frac{1}{32\pi^2} \ln \frac{M_\pi^2}{\mu^2} - \frac{1}{32\pi^2}\right) \frac{M_\pi^2}{F_\pi^2} + \dots \quad (12)$$

leading to the prediction[16]

$$\alpha = 1 + 0.06 + 0.01 + \dots = 1.07 \pm 0.007 . \quad (13)$$

(Here, the contributions of successive orders $O(p^2)$, $O(p^4)$ and $O(p^6)$ are explicitly shown).

This has to be compared with the experimental value[13]

$$\alpha_{exp} = 2.16 \pm 0.86 \quad (14)$$

inferred from the Geneva-Saclay K_{l4} [18] and CERN-Münich $\pi N \rightarrow \pi\pi N$ [17] data, using the technique of Roy dispersion relations. Notice that the result (14) implies for the $q\bar{q}$ condensate $2mB = (0.6 \pm 0.4)M_\pi^2$. Results of new high precision experiments, such as the $\pi^+\pi^-$ - atom lifetime measurement at CERN[8] and the new K_{l4}^+ experiments [9] at BNL and Frascati are awaited with an obvious interest.

5. THE STRANGE QUARK MASS

Treating also the s-quark as light, the $G\chi PT$ formula (9) can be extended to remaining unmixed Goldstone bosons :

$$\begin{aligned} M_\pi^2 &= 2mB_0 + 4m^2 A_0 + O(p^3) \\ M_K^2 &= (m_s + m)B_0 + (m_s + m)^2 A_0 + O(p^3) . \end{aligned} \quad (15)$$

The constants B_0 and A_0 differ from B and A introduced before by Zweig rule violating effects, which are expected to be small. Eq. (15) implies (the dots stand for higher order corrections)

$$\frac{2mB_0}{M_\pi^2} = \frac{(r - r_1)(r + r_1 + 2)}{r^2 - 1} + \dots \quad (16)$$

$$r = r_2 - 2A_0 \left(\frac{m_s}{M_\pi}\right)^2 + \dots , \quad (17)$$

where r is the quark mass ratio $r = m_s/m$ and

$$r_n = 2 \left(\frac{M_K}{M_\pi}\right)^n - 1 , \quad n = 1, 2 , \quad (18)$$

i.e. $r_1 \simeq 6.3$ and $r_2 \simeq 25.9$. Notice that, according to Eq. (16), the stability of the vacuum condition $B_0 \geq 0$, requires $r \geq r_1 \simeq 6.3 + \dots$. It is instructive to look at Eq. (17) in the light of the new **experimental** information on m_s , obtained by the ALEPH-collaboration from the precise measurement of the inclusive rate of the τ -decays $\tau \rightarrow \nu_\tau + X$ ($S = 1$). The preliminary result reads[19]

$$m_s(1 \text{ GeV}) = (235_{-42}^{+35}) \text{ MeV} \quad (19)$$

and it is somewhat higher than expected on the basis of sum rule[20] and lattice[21] estimates.

A higher value of m_s is welcome for the $G\chi PT$ scenario : It **lowers** the values of the quark mass ratio r and of the condensate parameter B_0 . To illustrate this point by a crude numerical estimate, let us take as the input (at the QCD scale $\nu = 1$ GeV), $m_s = 200$ MeV and $A_0 \simeq A \simeq 4$ according to Eq. (8). Eq. (17) then yields $r \simeq 8.3$, implying in turn, through Eq. (16), $2mB_0 \simeq 0.5 M_\pi^2$, in agreement with the available information from $\pi\pi$ -scattering. The condensate should then be $B_0 \simeq 190$ MeV and m (1 GeV) $\simeq 24$ MeV. Let us stress once again that these numbers constitute a crude estimates with neither error bars nor higher order effects included. A more detailed analysis will be given elsewhere[22].

6. $m_u + m_d$ AND QCD SUM RULES

We finally discuss the constraints imposed on the value of $m = \frac{1}{2}(m_u + m_d)$ by the QCD sum rules involving the two point function

$$\Psi_5(q^2) = i \int dx e^{iqx} \times \\ \times \langle \Omega | T D_5(x) D_5^+(0) | \Omega \rangle , \quad (20)$$

where D_5 stands for the divergence of the axial current

$$D_5 \equiv \partial^\mu \bar{u} \gamma_\mu \gamma_5 d = 2m \bar{u} i \gamma_5 d . \quad (21)$$

Evaluation of these sum rules needs information about the imaginary part

$$\frac{1}{\pi} \text{Im} \Psi_5(t) = 2F_\pi^2 M_\pi^4 \delta(t - M_\pi^2) + \rho(t) . \quad (22)$$

The spectral function $\rho(t)$ collects all the continuum contributions of 3π , $K\bar{K}\pi$, $5\pi \dots$ intermediate states. It is in principle measurable in high statistics τ -decays [23] but so far, both its shape and its **normalization** are unknown. $\rho(t)$ is a product of a genuine massless QCD correlation function (similar to the one in Eq. (6)) and of m^2 . Since, in QCD, m is a free parameter, there is no way how to deduce the normalization of ρ from the theory, without an independent experimental input.

Two devices have been used in the literature to circumvent this difficulty. The first one attempts to use the χPT expression of $\rho(t)$ near threshold

to guess its normalization in the resonance region [24]. It turns out that this χPT -based normalization is surprisingly sensitive to the size of the $q\bar{q}$ condensate. At **threshold**, one has up to the leading order [23,12]

$$\rho|_{B=0} = 13.5 \times \rho|_{S\chi PT} . \quad (23)$$

The $G\chi PT$ knows that a substantially larger value of the quark mass m is needed for its internal consistency.

The second attempt to constrain the normalization of ρ makes use of QCD-hadron duality [25] : It requires [24] that the ratio

$$R_{Had}(s) = \frac{3}{2s} \frac{\int_0^s dt t \rho(t)}{2F_\pi^2 M_\pi^4 + \int_0^s dt \rho(t)} \quad (24)$$

coincides in a suitable range of s with the corresponding QCD expression $R_{QCD}(s)$ which is given by OPE and is almost independent of m . For small spectral densities, R_{Had} is, indeed, rather sensitive to the normalization of ρ . However, to the extent that the continuum contribution in the denominator of (24) dominates over the pion contribution, Eq. $R_{Had} = R_{QCD}$ becomes independent of the normalization of ρ , constraining merely its shape. This is what is **expected** to happen as a consequence of the $G\chi PT$ power counting (4) : $\int dt \rho(t)$ counts as $O(p^2)$, whereas the pion contribution is $O(p^4)$. It is actually not difficult to construct a model of the shape of $\rho(t)$, for which the duality criterion would be satisfied [22] **without including the contribution of the pion pole**. After all, in the scalar channel there is no pion pole and the corresponding $R_{QCD}(s)$ are about the same in both pseudoscalar and scalar channels.

Even if the normalization of the spectral function remains unknown, the QCD sum rules can still be combined with the Ward identity

$$\Psi_5(0) = -2m \langle \Omega | \bar{u}u + \bar{d}d | \Omega \rangle \quad (25)$$

to investigate the variation of $\Psi_5(0)$ with the quark mass m , **keeping fixed** $F_\pi^2 M_\pi^2$ at its experimental value. Notice that (25) does not directly involve the condensate (2) which is defined in the chiral limit. For small m , however,

$$\frac{1}{F^2} \Psi_5(0) = 4mB + 2m^2C + \dots , \quad (26)$$

where C is a high energy counterterm which depends on the way one renormalises $\Psi_5(0)$. We now consider [22] the Laplace transform sum rule for $\Psi_5(-Q^2)/Q^2$ making appear $\Psi_5(0)$, together with its derivatives. We use a simple model for the shape of the spectral function :

$$\rho(t) = [m(1 \text{ GeV})]^2 \{g^2[\delta(t - M_1^2) + x\delta(t - M_2^2)] + \gamma_{as}(t)\theta(t - t_0)\} \quad (27)$$

where $M_1 = 1.30 \text{ GeV}$, $M_2 = 1.77 \text{ GeV}$, $x \simeq 1$ and $\gamma_{as}(t)$ is given by QCD asymptotics. Eliminating the unknown constant g^2 one obtains for each value of m a prediction for the condensate ratio $\Psi_5(0)/2F_\pi^2 M_\pi^2$, which in $S\chi PT$ is expected to be close to 1. The result is displayed in Fig. 1. The indicated errors reflect the variation of




Figure 1. The condensate ratio $\Psi_5(0)/2F_\pi^2 M_\pi^2$ as a function of $m(1\text{GeV})$ from Borel transform sum rules for $\Psi_5(q^2)$.

individual output values with the Borel variable u . The shape of the curve on Fig. 1 can be understood within $G\chi PT$: Using the expansion (5) and Eq. (26), one gets

$$-\frac{m}{F_\pi^2 M_\pi^2} < \Omega | \bar{u}u + \bar{d}d | \Omega > =$$

$$= 1 - 4 \frac{m^2}{M_\pi^2} (A - \frac{1}{4}C) + \dots \quad (28)$$

which reproduces the curve in the whole range of $m \lesssim 35 \text{ MeV}$, provided one takes

$$A - \frac{1}{4}C \simeq 4 \div 5.4 \quad (29)$$

(independent of m). It is instructive to compare this result with our estimate (8).

7. SUMMARY

The leading order $G\chi PT$ expression of M_π^2 and M_K^2 involves **i)** the condensate parameter B_0 , **ii)** the quadratic slope parameter A_0 , **iii)** $m = \frac{1}{2}(m_u + m_d)$ and **iv)** m_s . Consequently, in order to get a complete picture of quark masses and condensates, it is sufficient to pin down any two of these 4 parameters independently of any (hidden) prejudices about the size of the $q\bar{q}$ condensate. I have suggested to use recent experimental determination of m_s by the ALEPH collaboration, together with an estimate of A_0 . I have argued that A_0 should be roughly the same both in $S\chi PT$ and in $G\chi PT$, rather insensitive to the $q\bar{q}$ -condensation. The crude estimate $A_0 \simeq 4$ (with errors hard to determine) then leads to a rather small values of $r = m_s/m \simeq 8.3$ and of the condensate parameter $B \sim 190 \text{ MeV}$. $2mB$ then represents half of M_π^2 , in agreement with existing $\pi - \pi$ scattering data. The corresponding large value of $m \simeq 24 \text{ MeV}$ has been argued to be compatible with relevant QCD sum rules and with QCD-hadron duality, provided the hadronic spectral function is indeed larger than usually expected. This can be tested measuring azimuthal asymmetries in $\tau \rightarrow 3\pi + \nu_\tau$ [23].

The preceeding discussion certainly lacks a quantitative precision and it should be merely viewed as a plausibility argument illustrating the coherence of the small condensate scenario. The precise quantitative test is awaited in low-energy $\pi^+ \pi^-$ scattering.

I am indebted to Norman Fuchs, Marc Knecht and Bachir Moussallam for discussions and help in collecting materials for this talk.

REFERENCES

1. G. 't Hooft in : Recent development in gauge theories, eds. G. 't Hooft et al. (Plenum, New York, 1980) ; Y. Frishman et al., Nucl. Phys. B177 (1981) 157 ; S. Coleman and B. Grossman, Nucl. Phys. B203 (1982) 205.
2. C. Vafa and E. Witten, Nucl. Phys. B234 (1984) 173 ; Comm. Math. Phys. 95 (1984) 257.
3. T. Banks and A. Casher, Nucl. Phys. B168 (1980) 103 ; H. Leutwyler and A.V. Smilga, Phys. Rev. D46 (1992) 5607.
4. J. Stern, to be published.
5. R. Gupta in LATTICE 94, Nucl. Phys. B (Proc. Suppl.) 42 (1995) 85.
6. G. Arvanitis, F. Geniet, J.L. Kneur, A. Neveu, Phys. Lett. B390 (1997) 385 ; J.L. Kneur, hep-ph/96 09 265 and these proceedings.
7. M. Gell-Mann, R.J. Oakes and B. Renner, Phys. Rev. 175 (1968) 2195.
8. B. Adeva et al., preprint CERN/SPSLC 95-1, Dec 1994.
9. M. Baillargeon and J.P. Franzini, in "The second Daphne Physics Handbook" Eds. L. Maiani, N. Paver and G. Pancheri, INFN-LNF publication, May 1995. J. Lowe, private communication.
10. J. Gasser and H. Leutwyler, Ann. Phys. 158 (1984) 142 ; Nucl. Phys. B250 (1985) 465.
11. N.H. Fuchs, H. Sazdjian, J. Stern, Phys. Lett. B269 (1991) 183 ; J. Stern, H. Sazdjian, N.H. Fuchs, Phys. Rev. D47 (1993) 3814.
12. M. Knecht and J. Stern, in "The second Daphne Physics Handbook", Eds. L. Maiani, N. Paver and G. Pancheri, INFN-LNF publication, May 1995.
13. M. Knecht, B. Moussallam, J. Stern and N.H. Fuchs, Nucl. Phys. B457 (1995) 513 ; Nucl. Phys. B471 (1996) 445.
14. S. Weinberg, Phys. Rev. Lett. 17 (1966) 616 ; J. Gasser and H. Leutwyler, Phys. Lett. B125 (1983) 321.
15. J. Bijnens, G. Colangelo, G. Ecker, J. Gasser and M. Sainio, Phys. Lett. B374 (1996) 210.
16. L. Girlanda, M. Knecht, B. Moussallam and J. Stern, hep-ph/ 97 03 448, Phys. Lett. B in press.
17. B. Hyams et al., Nucl. Phys. B64 (1973) 134 ; G. Grayer et al., Nucl. Phys. B75 (1974) 189.
18. L. Rosselet et al., Phys. Rev. D15 (1977) 574.
19. S. Chen, these proceedings.
20. M. Jamin, these proceedings.
21. C. Allton, these proceedings ; H. Hoerber, these proceedings.
22. N.H. Fuchs, M. Knecht and J. Stern, to be published.
23. J. Stern, M. Knecht and N.H. Fuchs, in "Third Workshop on the tau-charm Factory", Marbella (1993), Eds. J. Kirkby and R. Kirkby, Editions Frontières 1994.
24. C.A. Dominguez and E. de Rafael, Ann. Phys. 174 (1987) 372 ; J. Bijnens, J. Prades and E. de Rafael, Phys. Lett. B348 (1995) 226.
25. J. Prades, these proceedings.